



ORTHENE® 97

(SOLUBLE INSECTICIDE)

Active Ingredient	By Wt.
Acephate (O,S-Dimethyl acetylphosphoramidothioate).....	97.4%
Other Ingredients.....	2.6%
Total.....	100.0%
U.S. Pat. No. 6,013,272	

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist. Wash hands thoroughly after handling. Do not allow children or pets to come into contact with treated areas until sprays have dried.

FIRST AID

Acephate is an organophosphate, cholinesterase inhibitor.

If swallowed: Call poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by the poison control center or doctor.

Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

Call a poison control center or doctor for treatment advice.

(continued)

FIRST AID (continued)

If inhaled: Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-892-0099** for emergency medical treatment information.

NOTES TO PHYSICIAN

This material contains a cholinesterase inhibitor. Measurement of blood cholinesterase activity may be useful in monitoring exposure but decisions regarding treatment will usually need to be made before test results are available. If signs of cholinesterase inhibition appear, atropine sulfate is antidotal. 2-PAM (PROTOPAM) is also antidotal and may be used in conjunction with atropine but should not be used alone.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to birds. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or soil-incorporate spills.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For non-agricultural areas, do not enter treated areas without protective clothing until sprays have dried.

DISCLAIMER, RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND LIMITATION OF LIABILITY

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is
(continued)

(continued)

reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the **Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

GENERAL INFORMATION

TABLE OF CONTENTS

BEANS AND LIMA BEANS –	
Dry and Succulent Forms.....	Table 1
CELERY.....	Table 2
COLE CROPS.....	Table 3
COTTON.....	Table 4
COTTON TANK MIXES.....	Table 5
COTTON TANK MIXES WITH	
PYRETHROIDS.....	Table 6
CRANBERRY.....	Table 7
HEAD LETTUCE – Crisphead Type Only.....	Table 8
MINT.....	Table 9
NON-BEARING CITRUS – RECOMMENDATIONS	
FOR USE IN AGRICULTURAL SETTINGS..	Table 10
NON-CROP AREAS – FIELD BORDERS,	
FENCEROWS, ROADSIDES, DITCHBANKS,	
BORROW PITS.....	Table 11
NON-CROP AREAS – WASTELAND.....	Table 12
NURSERY STOCK – NON-BEARING DECIDUOUS	
FRUIT TREES, NUT TREES AND VINES...	Table 13
PEANUTS.....	Table 14
PEPPERS – BELL.....	Table 15
PEPPERS – NON-BELL.....	Table 16
TOBACCO – TRANSPLANT WATER APPLICATION	
(EXCEPT CA).....	Table 17
TOBACCO – FOLIAR APPLICATION	
(EXCEPT CA).....	Table 18
TOBACCO – PLANT BED APPLICATION	
(EXCEPT CA).....	Table 19
TOBACCO – GREENHOUSE APPLICATION	
(EXCEPT CA).....	Table 20
TOBACCO – SOIL APPLICATION	
(EXCEPT CA).....	Table 21

DIRECTIONS

CHEMIGATION

Apply to Cranberries only by sprinkler irrigation systems. Do not apply by chemigation to any other crop, or this crop using any other type of irrigation system.

General Mixing Instructions

Thoroughly clean all sprayer components prior to mixing. Add approximately 1/2 of the required amount of water to the spray tank and begin agitation. Add the required amount of Orthene® 97 Soluble Insecticide and mix thoroughly. Then add other tank mix partners or foliar feed additives and the remaining water. Maintain agitation during filling and spraying to ensure a uniform spray mixture.

For application equipment, which has minimal agitation such as tobacco transplant water equipment, proper attention to mixing the *Orthene 97* product should be given. With tobacco transplant water applications the *Orthene 97* product should be premixed in water to form a slurry prior to putting the product into the transplant water applicator. If premixing is not done, then adequate time should be allowed for the product to dissolve in the transplant water prior to beginning application.

BEANS AND LIMA BEANS – Dry and Succulent Forms

TABLE 1. Beans and Lima Beans – Dry and Succulent Forms – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 20 to 100 gals./A of spray. By Air: Minimum of 2 gals./A of spray.	Fleahoppers Grasshoppers	1/4 to 1/2 lb. (4 to 8 oz.)	Apply when eggs or insects first appear. Repeat at 7 to 10 day spray intervals as necessary to maintain control.* Use higher rates for severe insect infestations.	14 (snapbeans – succulent or dry beans) 0 (lima beans – succulent form)
	Aphids (excluding Black Bean Aphid) Bean Leaf Beetle Bean Leafroller Cabbage Looper Cutworms Green Cloverworm Leafhoppers Mexican Bean Beetle Plantbugs (Lygus) Soybean Looper Thrips Whitefly (Except Sweetpotato or Silverleaf Whitefly)	1/2 to 1.0 lb. (8 to 16 oz.)		
	Armyworms (excluding Beet Armyworm) Corn Earworm European Corn Borer	3/4 to 1.0 lb. (12 to 16 oz.)		

USE PRECAUTIONS:

Do not feed treated vines to livestock.

*Do not apply more than 2-1/8 lbs./A (2 lbs. ai) per season.

CELERY

TABLE 2. Celery – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 50 to 100 gals./A of spray. By Air: Minimum of 5 gals./A of spray.	Green Peach Aphid	1/2 to 1.0 lb. (8 to 16 oz.)	Apply when eggs or insects first appear. Repeat at 3 to 10 day spray intervals as necessary to maintain control.*	21
	Cabbage Looper Fall Armyworms	1.0 lb. (16 oz.)		

USE PRECAUTIONS:

All celery must be trimmed (tops removed) before shipment for use.
 Do not use trimmed tops for food or feed.
 *Do not apply more than 2-1/8 lbs./A (2 lbs. ai) per season.

COLE CROPS

BRUSSELS SPROUTS & CAULIFLOWER

TABLE 3. Cole Crops – Brussels Sprouts & Cauliflower – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 25 to 150 gals./A of spray. By Air: Minimum of 5 gals./A of spray.	Green Peach Aphid	1/2 to 1.0 lb. (8 to 16 oz.)	Use the high rate when heavy infestations of aphids are present.	14
	Cabbage Looper Diamondback Moth Larvae Imported Cabbage Worm	1.0 lb. (16 oz.)	Apply when eggs or insects first appear. Repeat application as necessary to maintain control.* Diamondback Moth Larvae – This insect has demonstrated an ability to develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current recommended control practices for this insect.	

USE PRECAUTIONS:

*Do not apply more than 2-1/8 lbs./A (2 lbs. ai) per season to Brussels Sprouts and Cauliflower.
 Do not feed trimmings to livestock or allow animals to graze in treated areas.

COTTON

GENERAL USE PRECAUTIONS

Do not use treated seed for food or feed purposes or process for oil.

Do not feed gin trash to livestock.

Do not allow animals to graze on treated areas.

**Do not apply more than 6-1/8 lbs./A (6 lbs. ai) per season. This includes the use of *Orthene* in commercial seed treatment, in-furrow spray, in-furrow and foliar applications.

TABLE 4. Cotton – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
IN FURROW By Ground: 3 to 5 gals./A of spray.	Aphids* Black (Greasy) Cutworm (Except CA) Thrips (including Western Flower Thrips)	1/2 to 1.0 lb. (8 to 16 oz.) NOTE: For the Blacklands of Texas use 1.0 lb./A Orthene 97.	Use flat-fan nozzles for in-furrow application. Align nozzles to ensure good spray deposition into the seed furrow. Cone type nozzles are not recommended since unacceptable spray deposition may occur. Securely fasten spray-system tubing to furrow opener and check frequently to ensure proper positioning and operation. <i>Orthene 97</i> can be mixed with fungicides that are sprayed in-furrow for disease control.	
	*Excluding Cotton Aphids in AZ & CA			
FOLIAR By Ground: 10 to 25 gals./A of spray. By Air: 3 to 10 gals./A of spray (minimum of 5 gals./A in AZ & CA).	Thrips (including Western Flower Thrips)	2.5 oz. to 3.0 oz.	Apply when eggs or insects first appear. Repeat application as necessary to maintain control.**	21
	Plantbugs (Lygus)	1/4 to 1.0 lb. (4 to 16 oz.) 3/4 to 1.0 lb. (AZ & CA)	Apply when eggs or insects first appear. Use highest rate for Lygus adults that have migrated into cotton. Repeat application as necessary to maintain control.**	21
	Fleahopper	1/4 lb. (4 oz.)	Apply when eggs or insects first appear. Repeat application as necessary to maintain control.**	21
	Cotton Aphid (excluding AZ & CA)	1/2 to 1.0 lb. (8 to 16 oz.)	This insect may develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current control recommendations. Repeat application as necessary to maintain control.**	21
	Whitefly (excluding Sweetpotato Whitefly/Silverleaf Whitefly)	1/2 to 1.0 lb. (8 to 16 oz.)	Repeat application as necessary to maintain control.**	21
	Armyworms (excluding Beet Armyworm) Cabbage Looper	1.0 lb. (16 oz.)	Apply when eggs appear and repeat at 3 to 7 day spray intervals.**	21

(continued)

COTTON (continued)

TABLE 4. Cotton – Recommendations for Use (continued)

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 10 to 25 gals./A of spray. By Air: 3 to 10 gals./A of spray (minimum of 5 gals./A in AZ & CA).	Bollworm Tobacco Budworm Adults Larvae Eggs/Deadhatch®	1/2 to 1.0 lb. (8 to 16 oz.) (East of Rockies) 1.0 lb. (16 oz.) (AZ & CA)	Early Season: Use 1/2 lb. /A for light infestation. Mid and Late Season: Use 3/4 to 1.0 lb. /A for moderate to severe infestations. Apply when eggs appear and repeat at 3 to 7 day spray intervals.** Moths of budworm larvae are controlled by direct contact with spray. Moth kill is most likely to occur when late evening applications are made. Deadhatch: Control of emerging larvae by consumption of treated egg casings.	21
	Stinkbugs	3/4 lb. (12 oz.)	Apply when eggs appear and repeat at 3 to 7 day spray intervals.**	21
	Pink Bollworm (AZ & CA)	1.0 lb. (16 oz.)	Apply when insects appear and repeat at 5 to 7 day spray intervals.**	21
	Cutworms	3/4 lb. (12 oz.)	Ground application is recommended. Aerial applications are less effective, but may be used. Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant. Apply when insects first appear or damage is first noted and repeat application as necessary to maintain control.**	21

COTTON TANK MIXES

GENERAL USE PRECAUTIONS

Do not feed gin trash or treated forage to livestock.

Do not allow livestock to graze on treated areas.

*Do not apply more than 6-1/8 lbs./A (6 lbs. ai) per season. This includes the use of *Orthene* in commercial seed treatment, in-furrow spray, in-furrow and foliar applications.

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions which appear on all product labels. The most restrictive labeling applies when using a tank mix.

TABLE 5. Cotton Tank Mixes – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	ORTHENE 97 AND TANK MIX PARTNER	AMOUNT OF ORTHENE 97 + TANK MIX PARTNER PER ACRE	REMARKS
FOLIAR By Ground: 10 to 25 gals./A spray. By Air: 3 to 10 gals./A of spray (minimum 5 gals./A in CA).	Sweetpotato Whitefly (Silverleaf Whitefly)	<i>Orthene</i> 97 + Danitol® 2.4 EC	1/2 lb. (8 oz.) + 8 to 16 fl. oz.	User should comply with all applicable directions, restrictions and precautions on the EPA registered label for <i>Danitol</i> 2.4 EC Spray. Use a minimum application rate of 10-2/3 fl. oz./acre <i>Danitol</i> (0.2 lb. ai./acre) in CA and AZ.
	Armyworms (excluding Beet Armyworm) Bollworm Cabbage Looper Cotton Aphid Cutworms Fleahopper Grasshoppers Pink Bollworm Plantbugs (Lygus, Mirids) Salt Marsh Caterpillar Thrips (including Western Flower Thrips) Tobacco Budworm Whitefly (Excluding Sweetpotato/Silverleaf Whitefly)	<i>Orthene</i> 97 + Lorsban® 4E	1/4 to 1.0 lb. (4 to 16 oz.) + 1 to 2 pts.	User should comply with all applicable directions, restrictions and precautions on the EPA registered label for Lorsban 4E. Cutworms: Use 3/4 lb./A of <i>Orthene</i> 97. By Ground: Ground application is recommended. Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant. By Air: Aerial applications are less effective, but may be used. Apply when insects first appear or when damage is first noted and repeat application as necessary to maintain control.*

RESISTANCE-MANAGEMENT

Cotton pest control programs, especially those for control of Silverleaf Whitefly populations, should employ a properly designed resistance-management strategy. Such resistance-management strategies include mixture or rotation of alternative classes of chemistry including organophosphates, carbamates, pyrethroids or insect growth regulators. Consult your state or area agricultural extension service for local resistance-management strategies and advice on alternative insecticides.

COTTON TANK MIXES WITH PYRETHROIDS

GENERAL USE PRECAUTIONS

Do not feed gin trash or treated forage to livestock.

Do not allow livestock to graze on treated areas.

**Do not apply more than 6-1/8 lbs./A (6 lbs. ai) per season. This includes the use of *Orthene* in commercial seed treatment, in-furrow spray, in-furrow and foliar applications.

Synthetic Pyrethroids should be used within the guidelines of state and/or regional resistance management programs and recommendations.

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions which appear on all product labels. The most restrictive labeling applies when using a tank mix.

TABLE 6. Cotton Tank Mixes with Pyrethroids – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	ORTHENE 97 AND TANK MIX PARTNER	AMOUNT OF ORTHENE 97 + TANK MIX PARTNER PER ACRE	REMARKS
FOLIAR By Ground: 10 to 25 gals./A spray. By Air: 3 to 5 gals./A spray (minimum 5 gals./A in CA). See aerial remarks for Cutworms.	Aphids Bollworm Cabbage Looper Cotton Leaf Perforator Cutworms Fall Armyworm Fleahoppers Pink Bollworm (AZ & CA) Plantbugs Stinkbugs Sweetpotato/Silverleaf Whitefly Thrips (including Western Flower Thrips) Tobacco Budworm Whitefly	<i>Orthene 97</i>	1/2 to 1.0 lb./A (8 to 16 oz.)	Apply when eggs or insects first appear. Repeat application as necessary to maintain control.** Stinkbugs: Use 3/4 lb./A of <i>Orthene 97</i> . *Cutworms: Use 3/4 lb./A of <i>Orthene 97</i> . By Ground: Ground application is recommended. Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant. By Air: 3 to 10 gals./A spray (minimum 5 gals./A in CA) Aerial applications are less effective, but may be used. Apply when eggs or insects first appear or damage is first noted. Repeat application as necessary to maintain control.**
		+ one of the following:	+ one of the following:	
		Ammo [®] 2.5 EC*	Refer to the Ammo 2.5 EC approved label for use instructions.	
		Asana [®] XL*	Refer to the Asana XL approved label for use instructions.	
		Baythroid [®] 2 EC	Refer to the Baythroid 2 EC approved label for use instructions.	
		Capture [®] 2 EC	Refer to the Capture 2 EC approved label for use instructions.	
		Karate [®] 1 E (Except CA)	Refer to the Karate 1 E approved label for use instructions.	
Scout X-Tra [®] (Except CA)	Refer to the Scout X-Tra approved label for use instructions.			

RESISTANCE-MANAGEMENT

Cotton pest control programs, especially those for control of Silverleaf Whitefly populations, should employ a properly designed resistance-management strategy. Such resistance-management strategies include mixture or rotation of alternative classes of chemistry including organophosphates, carbamates, pyrethroids or insect growth regulators. Consult your state or area agricultural extension service for local resistance-management strategies and advice on alternative insecticides.

CRANBERRY

TABLE 7. Cranberry – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATE OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
<p>FOLIAR By Ground or Sprinkler: Sufficient water to give thorough coverage. By Air: Minimum of 2 gals./A of spray.</p>	<p>Cranberry Blossom Worm Cranberry Cutworm Gypsy Moth False Armyworm Fireworms Spanworms Sparganothis</p>	<p>1.0 lb. (16 oz.)</p>	<p>SPRINKLER IRRIGATION APPLICATION TO CRANBERRIES: This product may only be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.</p>	<p>90</p>
<p>USE PRECAUTIONS: Note: Limit to one application per growing season. Do not apply more than 1.0 lb. (1.0 lb. ai) per season. Do not apply from start of bloom until all berries set.</p>			<p>If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.</p>	
<p>Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment. Solid Set System: Apply specified dosage for the entire length of the irrigation period or for a 30 to 60 minute period at the end of a regular irrigation set or as a 30 to 60 minute injection as a separate application not associated with a regular irrigation. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining pesticide, a dye indicator may be injected into the lines to mark the end of the application period. See NOTE. Center Pivot Systems: Inject the specified dosage per acre continuously for one complete revolution of the system. See NOTE. NOTE: Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Inject the product with a positive displacement pump into the main line ahead of a right turn to ensure adequate mixing. Application of more than label-recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.</p>				

HEAD LETTUCE – Crisphead Type Only

TABLE 8. Head Lettuce (Crisphead Type Only) – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 10 to 60 gals./A of spray broadcast. By Air: Minimum of 5 gals./A of spray.	Aster Leafhopper Green Peach Aphid	1/2 to 1.0 lb. (8 to 16 oz.)	Repeat application as necessary to maintain control.*	21 Spring, Summer and early Fall crops in all areas, Winter crops in Florida and Texas, late Fall crops in Arizona and Winter crops in Arizona and California.
	Armyworms (excluding Beet Armyworm) Cabbage Looper	1.0 lb. (16 oz.)		

USE PRECAUTIONS:

*Do not apply more than 2-1/8 lbs./A (2 lbs. ai) per season.

Do not feed trimmings to livestock or allow animals to graze on treated areas.

AZ & CA, Desert Areas: Do not apply after first head begins to form in crops which germinate from mid-September through November.

MINT

PEPPERMINT AND SPEARMINT

TABLE 9. Mint – Peppermint and Spearmint – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 20 to 100 gals./A of spray. By Air: 5 to 10 gals./A of spray.	Alfalfa Looper Aphids Cutworms	1.0 lb. (16 oz.)	Apply when eggs or insects first appear. Make one repeat application, if necessary, to maintain control.*	14
FOLIAR By Ground: 40 to 100 gals./A of spray.	Strawberry Root Weevil Adult Black Vine Weevil Adult	1.0 lb. (16 oz.)	Strawberry Root and Black Vine Weevil: For control, apply in water in a minimum of 40 gals. spray per acre by ground. Good spray coverage and canopy spray penetration is critical for control. Increase spray volume and pressure with tall or dense mint canopy. Apply after adult emergence is complete but prior to egg laying. Apply at dusk or during the night on a warm still evening. Two applications 10 to 14 days apart may be necessary to reduce heavy infestations.	14

USE PRECAUTIONS FOR ALL APPLICATION METHODS:

*Do not apply more than 2-1/8 lb./A (2 lbs. ai) per season.

Do not graze treated areas.

**NON-BEARING CITRUS – RECOMMENDATIONS FOR USE IN AGRICULTURAL SETTINGS
(Newly Planted Groves)**

GENERAL USE PRECAUTIONS

Do not graze treated areas.
DO NOT HARVEST citrus for one year after treatment.

TABLE 10. Non-Bearing Citrus – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS
FOLIAR By Ground: 100 to 200 gals./A of spray.	Aphids Grasshoppers Katydid Mealybugs Orangedogs Plantbugs Thrips Whiteflies (except Sweetpotato/Silverleaf)	1/2 lb. (8 oz.)	Repeat application as necessary to maintain control.
	Citrus Blackfly*	1/2 to 3/4 lb. (8 to 12 oz.)	Apply when eggs or insects first appear. Use the high rate when a heavy infestation is present. Repeat at 7 to 10 day spray intervals.
	Ants Including Imported Fire Ants and Harvester Ants*	3/4 lb. (12 oz.)	Repeat application as necessary to maintain control.
Spray individual juvenile or non-bearing trees for coverage with total application not to exceed specified rate in lbs. per acre. Length of residual activity will depend upon spray coverage and the amount of moisture following application.			
SOIL MOUND TREATMENT DRENCH METHOD	Ants Including Imported Fire Ants and Harvester Ants*	3/4 oz./ 5 gals.	Apply 1 gal. of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound. Repeat application as necessary to maintain control.

*Not registered in CA for control of Citrus Blackfly and Harvester Ants.

NON-CROP AREAS

FIELD BORDERS, FENCEROWS, ROADSIDES, DITCHBANKS, BORROW PITS

TABLE 11. Non-Crop Areas – Field Borders, Fencerows, Roadsides, Ditchbanks, Borrow Pits – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATE OF ORTHENE 97 PER ACRE	REMARKS
FOLIAR By Ground: 10 to 20 gals./A of spray. By Air: 1 to 5 gals./A of spray (minimum of 5 gals./A in CA).	Grasshoppers	1/4 lb. (4 oz.)	For early to mid-season application. Use the higher volumes when spraying dense foliage. An approved drift retardant may be added to aid in controlling drift and reducing evaporation of aerial sprays.
USE PRECAUTION: Do not graze or feed vegetation cut from treated areas.			

(continued)

NON-CROP AREAS (continued)

WASTELAND

TABLE 12. Non-Crop Areas – Wasteland – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS
<p>FOLIAR</p> <p>By Ground: 10 to 20 gals./A of spray.</p> <p>By Air: 1/2 gal./A of spray (minimum of 5 gals./A in CA).</p>	<p>Black Grass Bugs Grasshoppers Mormon Crickets</p>	<p>1.5 oz. to 2 oz.</p>	<p>Use the higher volumes when spraying dense foliage.</p> <p>An approved drift retardant may be added to aid in controlling drift and reducing evaporation of aerial sprays.</p>
<p>USE PRECAUTION: Do not make more than one application per season. Do not graze or feed vegetation cut from treated areas.</p>			

NURSERY STOCK

NON-BEARING DECIDUOUS FRUIT TREES, NUT TREES AND VINES IN NURSERY FIELDS OR NON-BEARING ORCHARDS

TABLE 13. Nursery Stock – Non-Bearing Deciduous Fruit Trees, Nut Trees and Vines in Nursery Fields or Non-Bearing Orchards – Recommendations for Use

PLANTS	INSECTS	AMOUNT OF ORTHENE 97 PER ACRE	TIME OF APPLICATION	REMARKS
<p>Nursery Stock (Non-Bearing Deciduous Fruit Trees, Nut Trees, Vines) Almond Apricot Apple Cherry Grape Kiwi Pear Pistachio Plum Prune Walnut</p>	<p>Aphids Plantbugs (Lygus) Thrips</p>	<p>1/2 to 1 lb. (8 to 16 oz.)</p>	<p>Apply as needed for control of existing populations.</p>	<p>Apply <i>Orthene 97</i> in 50 to 200 gals. of water per acre by ground and not less than 5 gals. of water per acre by air.</p>
<p>USE PRECAUTIONS: For application only to non-bearing trees and vines in nursery fields or in non-bearing orchards. Do not apply to interplants or single trees in a bearing orchard. Do not harvest any fruit from sprayed trees for one year after application. Do not allow livestock to graze treated areas.</p>				

PEANUTS

TABLE 14. Peanuts – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATES OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 10 to 50 gals./A of spray. By Air: 5 to 10 gals./A of spray.	Grasshoppers	1/4 to 1/2 lb. (4 to 8 oz.)	Apply when eggs or insects first appear. Repeat application as necessary to maintain control.* <i>Orthene</i> can be tank mixed with registered at-cracking and early post-emergence peanut herbicides, provided those products do not prohibit tank mixes, provided the most restrictive of label limitations and precautions are observed, and provided no label dosage rates are exceeded. To determine physical compatibility, pour the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is considered physically compatible. When mixing wettable powder or dry flowable formulations, add and disperse these first, then add liquid pesticides. Conduct another compatibility test at concentrations which will be present in the irrigation lines. If there is any separation which cannot be remixed readily, Valent recommends that the combination not be used. Combinations should be kept agitated and should be applied immediately. Do not allow combinations to set for prolonged periods in the chemical supply tank or irrigation lines.	14 (of digging)
	Thrips	3/8 to 3/4 lb. (6 to 12 oz.)		
	Corn Earworm Fall Armyworm Leafhoppers Loopers Velvetbean Caterpillar	3/4 to 1.0 lb. (12 to 16 oz.)		
USE PRECAUTIONS: Do not feed treated forage or hay to livestock. Do not allow animals to graze on treated areas. *Do not apply more than 4-1/8 lbs./A (4 lbs. ai) per season.				

PEPPERS

BELL

TABLE 15. Peppers – Bell – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATE OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 25 to 150 gals./A of spray. By Air: Minimum of 3 gals./A of spray (minimum of 5 gals./A in CA).	Grasshoppers	1/4 to 1/2 lb. (4 to 8 oz.)	Apply when eggs or insects first appear. Repeat application as necessary to maintain insect-pest populations below economically damaging numbers.*	7
	Cabbage Looper Green Peach Aphid Tobacco Hornworm	1/2 to 1.0 lb. (8 to 16 oz.)		
	European Corn Borer	3/4 to 1.0 lb. (12 to 16 oz.)		

USE PRECAUTION:

*Do not apply more than 2-1/8 lbs./A (2 lbs. ai) per season.

NON-BELL

FOR USE IN MIDWESTERN, EASTERN STATES AND PUERTO RICO ONLY.

TABLE 16. Peppers – Non-Bell – Recommendations for Use

APPLICATION METHOD	PESTS CONTROLLED	RATE OF ORTHENE 97 PER ACRE	REMARKS	DAYS TO HARVEST
FOLIAR By Ground: 40 to 150 gals./A of spray.	Aphids	1/2 lb. (8 oz.)	Repeat at 7 to 10 day spray intervals as necessary.*	7

USE PRECAUTION:

*Do not apply more than 1.0 lb./A (1.0 lb. ai) per season.

TOBACCO

TOBACCO TRANSPLANT WATER APPLICATION (EXCEPT CA)

TABLE 17. Tobacco Transplant Water Application – Recommendations for Use

CROP	INSECTS	AMOUNT OF ORTHENE 97 PER ACRE	FURTHER USE INSTRUCTIONS
Tobacco	Cutworms Flea Beetle Green Peach Aphid Potato Tuberworm Tobacco Aphid Tobacco Thrips	3/4 lb. (12 oz.)	<p>Provides control of early season Flea Beetles, Green Peach Aphids, Tobacco Aphids, and Cutworms for approximately 3 to 4 weeks after transplanting. For later season control of these insects, apply a foliar spray of <i>Orthene 97</i>.</p> <p>Apply in a minimum of 100 gals. of transplant water per acre. <i>Orthene 97</i> should be pre-mixed in water to form a slurry solution prior to adding to the transplant water tank.</p> <p>Do not apply more than 3/4 lb. <i>Orthene 97</i> per acre as a transplant water application as some phytotoxicity may occur.</p> <p>Do not apply more than 4-1/8 lbs./A (4 lbs. ai) per season. This includes the use of <i>Orthene</i> in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.</p>

TOBACCO FOLIAR APPLICATION (EXCEPT CA)

TABLE 18. Tobacco Foliar Application – Recommendations for Use

CROP	INSECTS	AMOUNT OF ORTHENE 97 PER ACRE	FURTHER USE INSTRUCTIONS	DAYS TO HARVEST
Tobacco (Flue-Cured, Air Cured, Dark Fire Cured)	Grasshoppers	1/4 to 1/2 lb. (4 to 8 oz.)	Apply in 10 to 50 gals. water per acre with ground equipment or a minimum of 3 gals. per acre by air.	3
	Flea Beetle Green Peach Aphid Hornworm Tobacco Thrips	1/2 lb. (8 oz.)	Apply on a 7 day schedule or as needed.	
	Stinkbugs Tobacco Aphid Vegetable Weevils	1/2 to 3/4 lb. (8 to 12 oz.)	Do not apply more than 4-1/8 lbs./A (4 lbs. ai) per season. This includes the use of <i>Orthene</i> in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.	
	Budworm Cabbage Looper Cutworm Japanese Beetle	3/4 lb. (12 oz.)		

(continued)

TOBACCO (continued)

TOBACCO PLANT APPLICATION (EXCEPT CA)

TABLE 19. Tobacco Plant Bed Application – Recommendations for Use

CROP	INSECTS	AMOUNT OF ORTHENE 97 PER ACRE	FURTHER USE INSTRUCTIONS
Tobacco Bed Treatment	Cutworm Flea Beetle Green Peach Aphid Tobacco Aphid	3/4 lb. (12 oz.)	Apply to foliage at the equivalent of 3/4 table- spoon in 1 gal. of water per every 1,000 sq. ft. of bed. Apply evenly to insure thorough coverage. Do not apply more than 4-1/8 lbs./A (4 lbs. ai) per season. This includes the use of <i>Orthene</i> in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.

TOBACCO GREENHOUSE APPLICATION (EXCEPT CA)

TABLE 20. Tobacco Greenhouse Application – Recommendations for Use

CROP	INSECTS	AMOUNT OF ORTHENE 97 PER ACRE	FURTHER USE INSTRUCTIONS
Tobacco	Cutworm Flea Beetle Green Peach Aphid Tobacco Aphid	3/4 lb. (12 oz.)	Apply to foliage at the equivalent of 3/4 table- spoon in 3 gals. water per every 1,000 sq. ft. of bed. Apply evenly to insure thorough coverage. Note: Floatbed water should be disposed of in the transplanted field in either transplant water or foliar spray. Do not apply more than 4-1/8 lbs./A (4 lbs. ai) per season. This includes the use of <i>Orthene</i> in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.

TOBACCO SOIL APPLICATION (EXCEPT CA)

TABLE 21. Tobacco Soil Application – Recommendations for Use

CROP	INSECTS	AMOUNT OF ORTHENE 97 PER ACRE	FURTHER USE INSTRUCTIONS	DAYS TO HARVEST
Tobacco	Ants, Including Imported Fire Ants and Harvester Ants	Mound Treatment – Drench Method: Mix 3/4 oz. in 5 gals. of water. Apply 1 gal. of mix to each mound area by sprinkling the mound until it is wet and treat a four (4) foot diameter circle around the mound.	For best results apply the material in the early morning or late afternoon when the ants are most active. Applications made under pro- longed hot and dry conditions may be ineffective due to the location of the ants deep within the nest. Treat maximum of 13 mounds per acre. Do not treat more than once per season. Do not apply more than 4-1/8 lbs./A (4 lbs. ai) per season. This includes the use of <i>Orthene</i> in transplant water, plant bed, soil, float bed (greenhouse), and foliar applications.	3

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Open dumping is prohibited.

STORAGE

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers.

Store in cool, dry place. Protect from excessive heat.

Do not contaminate food or foodstuffs.

Do not store or transport near feed or food.

For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty canister into application equipment. Do not reuse container. Dispose of empty canister in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTE: This product is sold by weight and package is full when packed and is inclined to settle.

Eggs/Deadhatch is a registered trademark of Valent U.S.A. Corporation

Products That Work, From People Who Care is a trademark of Valent U.S.A. Corporation

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.

Orthene is a registered trademark of OMS Investments, Inc.

Ammo is a registered trademark of FMC Corporation

Asana is a registered trademark of E.I. du Pont de Nemours and Company

Baythroid is a registered trademark of Bayer Corporation

Capture is a registered trademark of FMC Corporation

Karate is a registered trademark of Syngenta

Lorsban is a registered trademark of Dow AgroSciences, LLC

Scout X-Tra is a registered trademark of Hoechst-Roussel Agri-Vet Co.

Manufactured for

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek CA 94596-8025

Made in U.S.A.

Form 1209-G

EPA Reg. No. 59639-91

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status or call 800-6-VALENT (682-5368).



For state registration and/or supplemental labels, please call or visit us online.

Products That Work, From People Who Care™ | www.valent.com | 800-6-VALENT (682-5368)

Read and follow the label instructions before using.

©2007 Valent U.S.A. Corporation. All rights reserved. Printed in the U.S.A.

2006-097-0001 5/07 IPCO/AV